Images in Clinical Tropical Medicine

Human Botfly Infestation in an Amerindian Girl in Southern Venezuela

Juan C. Gabaldon-Figueira,^{1*} Bobbi S. Pritt,² and Carlos Chaccour^{1,3,4}

¹Area de Enfermedades Infecciosas, Clínica Universidad de Navarra, Pamplona, Spain; ²Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, Minnesota; ³ISGlobal, Hospital Clínic, Universitat de Barcelona, Barcelona, Spain; ⁴Ifakara Health Institute, Ifakara, United Republic of Tanzania

A 9-year-old Pemón girl visited the primary healthcare outpatient clinic at Wonken, a Pemón village in the Gran Sabana region of Bolívar state, in southeastern Venezuela. Wonken is located within the Canaima National Park, a tropical rainforest region at 844 m above sea level. The average temperature is 20°C.

The patient presented a 4-week history of discomfort, itching, redness, and secretion in the back of her left thigh. She reported feeling movement inside the lesion, described as a “screwdriver being turned.” Despite living in the village, she indicated recent visits to Araparitá, a remote community south of Wonken.

She was afebrile and with stable vital signs. She presented a non-painful furuncle at the back of her left thigh, surrounded by an erythematous area of approximately 10 cm (Figure 1A). On close examination, a pair of moving respiratory spiracles were visible within the center of the lesion (Supplemental Video).

Suspecting a case of furuncular myiasis, a soap paste was applied on the lesion and covered in gauze to asphyxiate the fly larvae. After 3 hours, the gauze was removed, and the posterior end of the larva was clearly visible. A single live larva was successfully extracted (Figure 1B) and subsequently identified as a third instar stage larva (L3) of *Derma-*

tobia hominis, the human botfly.

The gross morphological characteristics that allowed for identification of the extracted larva included the cylindrical body shape with narrowed posterior end, and the backward-facing, coarse cuticular spines present on all but

the posterior three body segments (Figures 1C and D). The anterior mandibles (Figure 1E) were also clearly visible. Even though posterior spiracles were not examined (straight spiracular slits in a deep cavity are characteristic of this species), the gross morphologic appearance in the setting of furuncular myiasis in Venezuela is diagnostic of *D. hominis* infestation.¹

Received December 1, 2020. Accepted for publication December 15, 2020.

Published online February 22, 2021.

Note: Supplemental video appears at www.ajtmh.org.

Editor's Note: These images were previously published in *Dermatologia Venezolana*.

Authors' addresses: Juan C. Gabaldon-Figueira, Area de Enfermedades Infecciosas, Clínica Universidad de Navarra, Pamplona, Spain, E-mail: jgabaldonfi@unav.es. Bobbi S. Pritt, Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN, E-mail: pritt.bobbi@mayo.edu. Carlos Chaccour, Malaria Elimination, ISGlobal, Barcelona, Spain, and Malaria Elimination, Ifakara Health Institute, Morogoro, Tanzania, E-mail: carlos.chaccour@isglobal.org.

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC-BY) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

REFERENCE

1. Mathison BA, Pritt BS, 2014. Laboratory identification of arthropod ectoparasites. *Clin Microbiol Rev* 27: 48–67.

*Address correspondence to Juan C. Gabaldon-Figueira, Area de Enfermedades Infecciosas, Clínica Universidad de Navarra, Av. de Pío XII, 36, 31008 Pamplona, Navarra, Spain. E-mail: jgabaldonfi@unav.es

FIGURE 1. Case of *Dermatobia hominis* infestation. (A) Furuncle with visible respiratory spiracles. (B) The larva's body and posterior end are visible after occlusion of the lesion with soap paste. (C) Extracted L3 larvae with characteristic cylindrical body and narrowed posterior end. (D and E): Backward-facing cuticular spines and mandibles are visible on microscopic examination. This figure appears in color at www.ajtmh.org.